arXiv:gr-qc/9503045v1 24 Mar 1995

ON A HARMONIC PROPERTY OF THE EINSTEIN MANIFOLD CURVATURE

BABOUROVA O.V. and FROLOV B.N.

Department of Mathematics, Moscow State Pedagogical University,
Krasnoprudnaya 14, Moscow 107140, Russia;
E-mail: baburova.physics@mpgu.msk.su, frolovbn.physics@mpgu.msk.su

Abstract

The harmonicity condition of the curvature 2-form of a pseudo-Riemannian manifold is formulated on the basis of annulment of this form by the de Rham-Lichnerowicz Laplacian. The following theorem is proved: The curvature 2-form of any Einstein manifold is harmonic.

It is well known [1] that on the metric space-time manifold $\mathcal{M}$ the Maxwell equations can be expressed as

$$dF = 0\ , \qquad \delta F = 0\ , \tag{1}$$

where F is electromagnatic field strength 2-form and d, δ are the operations of exterior differentiation and codifferentiation, respectively. One has $\delta = *d*$ for even $n =$dim$\mathcal{M}$ with additional factor (-1) that occurs for space with negative signature. On the manifold $\mathcal{M}$ the Hodge-de Rham Laplacian $\Delta^{(HR)} = d\delta + \delta d$ can be constructed. The Maxwell equations (1) yield $\Delta^{(HR)}F = 0$, so the Maxwell 2-form F is *harmonic* [1]. The question is: whether the same result is valid for General Relativity, and if "yes" then in what sense.

On any Riemannian (pseudo-Riemannian) manifold one has the tensor-valued curvature 2-form

$$\Omega = \frac{1}{2}(R^{\lambda}{}_{\sigma\mu\nu} e_{\lambda} \otimes e^{\sigma}) \otimes \theta^{\mu} \wedge \theta^{\nu} \tag{2}$$

and the operation of covariant exterior differentiation $\mathcal{D}$ [2]. One can introduce the generalized operation of the covariant exterior codifferentiation as $\mathcal{D}^* = *\mathcal{D}*$ and *the de Rham-Lichnerowicz Laplacian*

$$\Delta^{(RL)} = \mathcal{D}\mathcal{D}^* + \mathcal{D}^*\mathcal{D} \ . \tag{3}$$

The operators $\mathcal{D}$, $\mathcal{D}^*$ and $\Delta^{(RL)}$ act not only on usual $\mathbf{R}$-valued forms but also on $\mathbf{V}$-valued forms (with values in a linear vector space $\mathbf{V}$).

We shall denote Ricci tensor components in local coordinate frames as $R_{\mu\nu} = R^{\lambda}{}_{\mu\lambda\nu}$. Then its trace $R = R^{\nu}{}_{\nu}$ is the curvature scalar. A Riemannian (pseudo-Riemannian) manifold with Ricci tensor satisfying the condition

$$R_{\mu\nu} = \Lambda g_{\mu\nu} \ , \qquad \Lambda = const \tag{4}$$

is called an Einstein manifold [3].

Definition: The curvature 2-form of Riemannian (pseudo-Riemannian) manifold is called *harmonic*, if one has

$$\Delta^{(RL)}\Omega = 0 \ . \tag{5}$$

We shall prove the following theorem.

Theorem: *The curvature 2-form of any Einstein manifold is harmonic.*

The proof of the theorem will be realized in a local coordinate frame $\theta^{\mu} = dx^{\mu}$. By generalizing the method developed in [4] to any (1,1)-tensor-valued 2-form Ω one has

$$\mathcal{D}\Omega = \frac{1}{3!}(3\Omega^{\lambda}{}_{\sigma\mu\nu;\rho} e_{\lambda} \otimes e^{\sigma}) \otimes \theta^{\rho} \wedge \theta^{\mu} \wedge \theta^{\nu} \ , \tag{6}$$

$$\mathcal{D}^*\Omega = (\Omega^{\lambda}{}_{\sigma\rho\nu}{}^{;\rho} e_\lambda \otimes e^\sigma) \otimes \theta^\nu \ , \tag{7}$$

$$\Delta^{(RL)}\Omega = \frac{1}{2}(\Omega^{\lambda}{}_{\sigma\mu\nu;\rho}{}^{;\rho} + 2\Omega^{\lambda}{}_{\sigma\mu}{}^{\rho}{}_{;[\rho;\nu]} - 2\Omega^{\lambda}{}_{\sigma\nu}{}^{\rho}{}_{;[\rho;\mu]}) e_\lambda \otimes e^\sigma \otimes \theta^\mu \wedge \theta^\nu \ . \tag{8}$$

Here $\Omega^{\lambda}{}_{\sigma\mu\nu;\rho}$ denotes the covariant derivative components with respect to all idices λ, σ, μ, ν. Applying now equation (8) to the curvature 2-form (2) and using the Ricci identity, one gets

$$\Delta^{(RL)}\Omega = \frac{1}{2}(\Delta^{(RL)}\Omega)^{\lambda}{}_{\sigma\mu\nu} e_\lambda \otimes e^\sigma \otimes \theta^\mu \wedge \theta^\nu \ , \tag{9}$$

$$\begin{aligned}(\Delta^{(RL)}\Omega)^{\lambda}{}_{\sigma\mu\nu} = {} & (R^{\lambda}{}_{\sigma\mu\nu;\rho}{}^{;\rho} + R^{\lambda}{}_{\sigma\nu}{}^{\alpha} R_{\alpha\mu} - R^{\lambda}{}_{\sigma\mu}{}^{\alpha} R_{\alpha\nu} \\ & -2R^{\lambda}{}_{\sigma\rho}{}^{\alpha} R_{\alpha\nu\mu}{}^{\rho} + 2R^{\lambda}{}_{\alpha\mu}{}^{\rho} R^{\alpha}{}_{\sigma\rho\nu} - 2R^{\lambda}{}_{\alpha\nu}{}^{\rho} R^{\alpha}{}_{\sigma\rho\mu}) \ . \end{aligned} \tag{10}$$

On the other hand, by covariant differentiating the Bianchi identity with the following contraction one gets

$$R^{\lambda}{}_{\sigma\mu\nu;\rho}{}^{;\rho} = R^{\lambda}{}_{\sigma\mu}{}^{\rho}{}_{;\nu;\rho} - R^{\lambda}{}_{\sigma\nu}{}^{\rho}{}_{;\mu;\rho} \ . \tag{11}$$

With the help of the Ricci identity (11) yields

$$\begin{aligned} R^{\lambda}{}_{\sigma\mu\nu;\rho}{}^{;\rho} = {} & R^{\lambda}{}_{\sigma\mu}{}^{\rho}{}_{;\rho;\nu} - R^{\lambda}{}_{\sigma\nu}{}^{\rho}{}_{;\rho;\mu} + R^{\lambda}{}_{\sigma\mu}{}^{\alpha} R_{\alpha\nu} - R^{\lambda}{}_{\sigma\nu}{}^{\alpha} R_{\alpha\mu} \\ & -2R^{\lambda}{}_{\sigma\rho}{}^{\alpha} R_{\alpha\mu\nu}{}^{\rho} + 2R^{\lambda}{}_{\alpha\mu}{}^{\rho} R^{\alpha}{}_{\sigma\nu\rho} - 2R^{\lambda}{}_{\alpha\nu}{}^{\rho} R^{\alpha}{}_{\sigma\mu\rho} \ . \end{aligned} \tag{12}$$

Now taking into account the following consequence of the Bianchi identity

$$R_{\mu\nu\sigma}{}^{\rho}{}_{;\rho} = 2R_{\sigma[\mu;\nu]} \ , \tag{13}$$

let us substitute (12) into the expression (10). Then we obtain the resultant expression for the action of the de Rham-Lichnerowicz Laplacian to the pseudo-Riemannian curvature 2-form:

$$\Delta^{(RL)}\Omega = 2R_{\mu[\lambda;\sigma]\nu} e^\lambda \otimes e^\sigma \otimes \theta^\mu \wedge \theta^\nu \ . \tag{14}$$

The expression (14) vanishes when the condition (4) is fulfilled. Therefore the curvature 2-form is harmonic when the given pseudo-Riemannian manifold is the Einsteinian one, as was to be proved.

It is easy to see that the converse statement is incorrect. There exist curvature 2-forms that are harmonic, but not Einsteinian.

The theorem proved above shows that the field strength harmonicity requirement is the reasonable condition also in the gravitatonal theory. This fact can be applied to the problem of Lagrangian choice in modern theories of gravitation, imposing the requirement of harmonicity on the corresponding curvature 2-forms.

References

1. T. Eguchi, P.B. Gilkey and A.J. Hanson, *Physics Reports* **66** (1980) 213.

2. R. Sulanke und P. Wintgen, *Differentialgeometrie und faserbündel* (Hochschulbücher für mathematik, band 75) (VEB Deutscher Verlag der Wissenshaften, Berlin, 1972).

3. A.L. Besse, *Einstein Manifolds* (Springer-Verlag, Berlin, Heidelderg, 1987).

4. G. de Rham, *Variétés Différentiables* (Hermann & C^{le}, Paris, 1955).